

634 287

SINGLE PULSE RESPONSE OF A PLASMA

Roy W. Gould

Technical Report No. 31

May 1966

CALIFORNIA INSTITUTE OF TECHNOLOGY

PASADENA, CALIFORNIA

SINGLE PULSE RESPONSE OF A PLASMA

Roy W. Gould

Technical Report No. 31

May 1966

Research sponsored by the
U. S. Office of Naval Research
Contract Nonr 220(50)

CALIFORNIA INSTITUTE OF TECHNOLOGY
Pasadena, California

SINGLE PULSE RESPONSE OF A PLASMA

Roy W. Gould

California Institute of Technology

ABSTRACT

The conjecture by Hirschfield and Wachtel¹ that a weak pulse of radiation can trigger the emission of a large delayed burst of coherent electron cyclotron radiation is examined critically. The effect is found to exist when the electrons have a single velocity perpendicular to the magnetic field, but disappears in a thermal equilibrium distribution.

In attempting to understand the cyclotron resonance echoes observed by Hill and Kaplan², Hirschfield and Wachtel have suggested a very interesting single pulse mechanism which results in large coherent delayed bursts of electron cyclotron radiation. The mechanism requires an extremely homogeneous magnetic field and an energy-dependent cyclotron frequency, such as that provided by the relativistic mass change. For electrons with a single velocity perpendicular to the field, v_0 , the current developed in plasma at the time of the first (of the many) delayed response was $|J| = .58 N e v_0$ where N is the number of electrons. This current is independent of the electric field required to trigger the burst, but the time at which the burst occurs is inversely proportional to the trigger field amplitude¹. Hirschfield and Wachtel assert that a similar effect occurs when the electrons have a Maxwellian velocity distribution except that only the first of the multiple bursts survives. In this note we rederive their result for monoenergetic electrons, then show that, upon integration over a Maxwellian velocity distribution, the effect disappears. Only a monotonically decaying response to the trigger pulse, which is proportional to the trigger pulse amplitude, survives.

It is convenient to describe the time-development of the electron velocities in a rotating velocity space which rotates with angular velocity ω of the applied electric field^{3,4}. The electric field is simply constant and produces a translation $v_1 = \frac{eE}{2m} t_{\text{pulse}}$ of each electron. We may neglect relativistic effects during the pulse provided that $v_1^2 \ll c^2$ and that the relativistic change in phase during the pulse $(\Delta\omega)_{\text{rel.}} t_{\text{pulse}}$, is negligible. Relativistic phase changes

following the pulse can be substantial even though the relativistic change in cyclotron frequency is small, simply because the time involved may be large. Prior to the pulse $\check{v} = v_{\perp} e^{i\phi_0}$, where v_{\perp} and ϕ_0 are the electron's initial speed and phase, respectively. The real and imaginary parts of \check{v} give v_x and v_y , respectively, in the rotating velocity space. Immediately following the pulse $\check{v} = v_{\perp} e^{i\phi_0} + v_{\parallel}$. In the time interval t following the pulse the electron rotates through an angle

$$\phi \cong \omega_{co} t \left(1 - \frac{v^2}{2c^2}\right) - \omega t \quad (1)$$

where $v^2 = v_{\perp}^2 + 2v_0 v_{\parallel} \cos \phi_0 + v_{\perp}^2 + v_{\parallel}^2$.

The square of the electron speed is assumed small compared to c^2 and only the leading term in the relativistic mass change is employed. The (complex) current at time t is given by

$$\check{J} = -e \int dn [v_{\perp} e^{i\phi_0} + v_{\parallel}] e^{i\phi} \quad (2)$$

where the integral is over the distribution of electrons.

We consider two cases: a) monoenergetic electrons, random phases, $dn = (N/2\pi) \delta(v_{\perp} - v_0) \delta(v_{\parallel}) dv_{\perp} dv_{\parallel} d\phi_0$, and b) Maxwell distribution of electron velocities, $dn = N(\pi v_{th}^2)^{-3/2} e^{-(v_{\perp}^2 + v_{\parallel}^2)/v_{th}^2} v dv_{\perp} dv_{\parallel} d\phi_0$.

For monoenergetic electrons (2) becomes

$$\check{J} = -Ne e^{i\omega' t} \int_0^{2\pi} \left[v_0 e^{i\phi_0} + v_{\parallel} \right] e^{-i\omega_{co} t \frac{v_0 v_{\parallel}}{c^2} \cos \phi_0} \frac{d\phi_0}{2\pi}$$

where $\omega' = \omega_{co} \left[1 - \frac{v_0^2 + v_{\parallel}^2}{2c^2} \right] - \omega$.

Using the identity $e^{-i\alpha \cos \phi_o} = \sum_{-\infty}^{\infty} (-i)^n J_n(\alpha) e^{-in\phi_o}$ and performing the ϕ_o integration yields

$$J = -Ne e^{i\omega'_c t} \left[-iv_o J_1(\omega_{co} t \frac{v_o v_1}{c^2}) + v_1 J_o(\omega_{co} t \frac{v_o v_1}{c^2}) \right] \quad (3)$$

This is the result of Hirschfield and Wachtel¹. When $v_1 \ll v_o$ the first term is generally more important [except at times when $J_1(\omega_{co} t \frac{v_o v_1}{c^2})$ is zero], the main effect of the small second term being to produce a small change in the zero-crossing times of the quantity in brackets. The important result is that the magnitude of the current exhibits periodic maxima, the first occurring when $\omega_{co} t \frac{v_o v_1}{c^2} = 1.84$ with a current $|J| = .58 Ne v_o$. The maximum current corresponds to a very strong phase bunching and is independent of v_1 , although the time required for the maximum current to develop increases as v_1 is decreased. This result indicates that a very weak pulse could trigger the emission of a very intense burst of cyclotron radiation in which most of the particles radiate coherently.

Hirschfield and Wachtel¹ have indicated that the main change introduced by assuming a Maxwellian distribution of electron velocities is to eliminate the secondary maxima, but that the first maximum remains and its characteristics may be estimated by using v_{th} , the electron thermal speed, for v_o in the above formulas. While at first sight this modification has an element of plausibility, the end result, that a weak trigger pulse can cause a nearly complete phase bunching of an equilibrium plasma, appears to violate the second law of thermodynamics. We now show that this

result is not correct.

When we assume dn to be given by a Maxwellian distribution [assumption (b)] we obtain

$$\begin{aligned} \tilde{J} = -Ne v_{th} I_{\parallel}(t) \int \frac{2v_{\perp} dv_{\perp}}{v_{th}^2} e^{-\frac{v_{\perp}^2}{v_{th}^2} - \frac{i\omega_{co} t}{2c^2} v_{\perp}^2} \left[-i \frac{v_{\perp}}{v_{th}} J_1(\omega_{co} t \frac{v_{\perp} v_1}{c^2}) + \right. \\ \left. + \frac{v_1}{v_{th}} J_0(\omega_{co} t \frac{v_{\perp} v_1}{c^2}) \right] \end{aligned} \quad (4)$$

$$\text{where } I_{\parallel}(t) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} e^{-\frac{v_{\parallel}^2}{v_{th}^2} - \frac{i\omega_{co} t}{2c^2} v_{\parallel}^2} \frac{dv_{\parallel}}{v_{th}} = \frac{1}{\sqrt{1 + i \frac{\omega_{co} t v_{th}^2}{2c^2}}} \quad (5)$$

comes from the integration over the component of velocity parallel to the magnetic field. The integral over perpendicular speeds can be accomplished with the integral

$$\left[\int_0^{\infty} J_{\nu}(ax) e^{-p^2 x^2} x^{\nu+1} dx = a^{\nu} e^{a^2/4p^2} / (2p^2)^{\nu+1} \right]$$

and yields:

$$\tilde{J}/(-Ne v_{th}) = \left[\frac{1}{\sqrt{1 + i\tau}} \right] \left[\frac{V_1}{1 + i\tau} - \frac{i\tau V_1}{(1 + i\tau)^2} \right] e^{-\frac{V_1^2}{2} \tau^2 / (1 + i\tau)} \quad (6)$$

where the current is expressed in units of $Ne v_{th}$, the "thermal current", $\tau = \omega_{co} t v_{th}^2 / c^2$ is the time expressed in units of the coherence time of cyclotron orbits, and $V_1 = v_1 / v_{th}$ is the strength of the "trigger" pulse expressed in units of the thermal speed. The first factor arises from the

integration over parallel velocities and contributes a decay $\sim \tau^{-1/2}$ when $\tau \gg 1$. The first and second terms in the second bracket arise from the integration of the Bessel functions J_0 and J_1 in (3) respectively. The important thing to note is that both contributions are now proportional to V_1 and that they combine in a simple way. Thus the plasma current induced by the pulse is strictly proportional to the amplitude of the pulse and furthermore the magnitude of the current decays monotonically from the initial value caused by the pulse:

$$\left| \frac{J}{N_e v_{th}} \right| = \frac{V_1}{(1 + \tau^2)^{5/4}} e^{-V_1^2 \tau^2 / (1 + \tau^2)} \quad (7)$$

When $V_1^2 \ll 1$ (weak pulse), the decay is governed by the first algebraic factor and the decay time is therefore

$$t_{decay} \approx \frac{1}{\omega_{co}} \frac{2c^2}{v_{th}^2}$$

The coherence time for cyclotron orbits (dephasing) arises because the cyclotron frequency depends slightly on particle energy and there is a distribution of energies.

One could therefore say that Hirschfield and Wachtel's mechanism fails to exist in a thermal plasma because the time required for phase bunching by their process exceeds the time for thermal dephasing by a factor $v_{th}/V_1 \gg 1$, and the latter process dominates. As a qualitative extension of this notion, one can also say that for the effect to exist in an almost monoenergetic distribution of perpendicular energies, such as in the experiment by Wachtel and Hirschfield⁵, the bunching time must be less than the

thermal dephasing time corresponding to the spread in velocities. This in turn sets a lower limit on the pulse strength, i.e., $v_{\perp} > \Delta v_{\perp}$, where Δv_{\perp} is a measure of the spread in perpendicular velocities.

In a similar fashion we can obtain a criterion for the required homogeneity of the magnetic field. When there is a spread in cyclotron frequencies of particles in the region of observation, this gives rise to an additional dephasing. In order that the one pulse maximum not be eliminated by magnetic field inhomogeneities $\Delta\omega_c \ll 1/t_{\max}$, or

$$\Delta\omega_{co} \ll \omega_{co} \frac{v_o v_{\perp}}{c^2} \quad (8)$$

where $\Delta\omega_{co}$ denotes the spread in cyclotron frequencies due to magnetic field inhomogeneities. This expression is also seen to imply a lower limit for v_{\perp} , for a given $\Delta\omega_{co}$.

The author acknowledges with pleasure numerous discussions with Dr. Wilhelm Kegel on this subject, and the support of the Office of Naval Research for research in this area.

REFERENCES

1. J. L. Hirschfield and J. M. Wachtel, 7th Ann. Am. Phys. Soc. Meeting Div. Plasma Physics 8-11 November 1965, Paper A-2, Bull. Am. Phys. Soc. (to be published).
2. R. M. Hill and D. E. Kaplan, Phys. Rev. Letters 14, 1062 (1965).
3. Roy W. Gould, Phys. Letters 19, 477 (1965).
4. Roy W. Gould, Bull. Am. Phys. Soc. 10, 1191 (1965); and Technical Report No. 28, Nonr 220(50), Dec. 1965.
5. J. M. Wachtel and J. L. Hirschfield, Bull. Am. Phys. Soc. 11, 375 (1966).

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION	
California Institute of Technology Pasadena, California			
		2b. GROUP	
3. REPORT TITLE			
SINGLE PULSE RESPONSE OF A PLASMA			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
Technical Report			
5. AUTHOR(S) (Last name, first name, initial)			
GOULD, Roy W.			
6. REPORT DATE		7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
May 31, 1966		7	5
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S)	
Nonr 220(50)		Technical Report No. 31	
b. PROJECT NO.			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.			
10. AVAILABILITY/LIMITATION NOTICES			
Distribution of this report is unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
		Office of Naval Research Washington, D. C.	
13. ABSTRACT			
<p>The conjecture by Hirschfield and Wachtel that a weak pulse of radiation can trigger the emission of a large delayed burst of coherent electron cyclotron radiation is examined critically. The effect is found to exist when the electrons have a single velocity perpendicular to the magnetic field, but disappears in a thermal equilibrium distribution.</p>			

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
coherent cyclotron radiation plasma emission pulse response of a plasma						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.